

Research Note

Helminths of the Western Lesser Siren, *Siren intermedia nettingi* (Caudata: Sirenidae), from Arkansas

CHRIS T. McALLISTER,¹ STEPHEN R. GOLDBERG,² STANLEY E. TRAUTH,³ CHARLES R. BURSEY,⁴ H. J. HOLSHUH,⁵ AND BETTY G. COCHRAN⁶

¹ Renal-Metabolic Lab (151-G), Department of Veterans Affairs Medical Center, 4500 S. Lancaster Road, Dallas, Texas 75216,

² Department of Biology, Whittier College, Whittier, California 90608,

³ Department of Biological Sciences, Arkansas State University, State University, Arkansas 72467,

⁴ Department of Biology, Pennsylvania State University, Shenango Valley Campus, 147 Shenango Avenue, Sharon, Pennsylvania 16146,

⁵ Comparative Medical and Veterinary Services, County of Los Angeles, Laboratory and Diseases Investigation, 12824 Erickson Avenue, Downey, California 90242 and

⁶ Caddo Ranger District, Ouachita National Forest, Glenwood, Arkansas 71943

ABSTRACT: Sixteen juvenile and adult western lesser sirens, *Siren intermedia nettingi* Goin, 1942, were collected from 2 counties in Arkansas and examined for endoparasites. Eleven (69%) sirens harbored 1 or more parasites, including 1 (6%) with *Diplostomum* sp. metacercariae, 9 (56%) with *Proteocephalus sireni*, 1 (6%) with larval *Capillaria* sp., and 2 (13%) with *Falcaustra chabaudi*. New locality records are documented for *P. sireni* and *F. chabaudi*, and new host records are reported for *Diplostomum* sp. and *Capillaria* sp. The histopathology of the larval *Capillaria* integumental infection is described in the new host. A summary is provided on helminths of *Siren* spp.

KEY WORDS: Amphibia, Arkansas, *Capillaria* sp., Caudata, *Diplostomum* sp., *Falcaustra chabaudi*, histopathology, intensity, prevalence, *Proteocephalus sireni*, Sirenidae, *Siren intermedia nettingi*, survey.

The western lesser siren, *Siren intermedia nettingi* Goin, 1942, is a large eel-like amphibian that ranges through the Mississippi River Valley east to western Alabama and west to eastern Texas (Martof, 1973; Conant and Collins, 1991). Much is known about the natural history and ecology of this salamander (Martof, 1973), including information on its helminths (Nickol, 1972; Dunagan and Miller, 1973; Dyer, 1973; Brooks and Buckner, 1976; Brooks, 1978; Buckner and Nickol, 1979). Except for Louisiana populations, these reports concern sirens from more northern parts of the range in Illinois. Herein, we report on helminths of a small sample of *S. i. nettingi* from Arkansas and provide a summary of helminths of North American *Siren* spp.

A total of 16 juvenile and adult specimens of *S. i. nettingi* (mean \pm SE snout-vent length [SVL] = 180.2 ± 85 , range 53–290 mm) were collected alive between October 1990 and March 1991, and again during January 1993, with minnow

traps, dip nets, or by hand at wetland sites in Montgomery ($N = 13$) and Clay ($N = 3$) counties of Arkansas and examined for endoparasites. Methods for necropsy, coccidial isolation, and preparation and staining of blood films and helminths follow those used by McAllister and Upton (1987). Voucher specimens of hosts are deposited in the Arkansas State University Museum of Zoology (ASUMZ). Specimens of parasites are deposited in the U.S. National Parasite Collection, Beltsville, Maryland 20705, as follows: *Diplostomum* sp. (USNM 82843), *Proteocephalus sireni* (USNM 82844), *Capillaria* sp. (USNM 82845), *Falcaustra chabaudi* (USNM 82842).

Eleven sirens (mean \pm SE SVL = 189.7 ± 20.3 , range 53–290 mm) were infected with 1 or more helminths, including *Diplostomum* sp. metacercariae, *Proteocephalus sireni*, larval *Capillaria* sp., and *Falcaustra chabaudi*. Two sirens harbored multiple infections of either larval *Capillaria* sp. and *P. sireni* or *Diplostomum* sp. metacercariae and *P. sireni*. Sirens were negative for coccidians, blood hematozoans, and myxozoans.

Numerous strigeoid metacercariae of the diplostomulum type were free in the coelomic and pericardial cavities of a single *S. i. nettingi* (SVL = 252 mm) collected in February 1991 from Montgomery County. The diplostomula appeared identical to the description of the metacercarial stage of *Diplostomum variabile* (Chandler, 1932) Dubois, 1937 (= *Didelphodiplostomum variabile* (Chandler, 1932) Dubois, 1944 (see Harris et al., 1967)). This present finding represents a new host record for *Diplostomum* sp.

A total of 39 tapeworms fitting the description of *Proteocephalus sireni* (Brooks and Buckner,


Table 1. Summary of helminths of *Siren* spp.

Host taxon/parasite	Locality	Reference(s)
<i>Siren intermedia nettingi</i>		
Trematoda		
<i>Allassosiomoides louisianensis</i>	Illinois	Brooks and Buckner, 1976
<i>Diplostomum</i> sp.	Arkansas	This report
<i>Progorogodera foliata</i>	Illinois	Brooks and Buckner, 1976
Cestoidea		
<i>Proteocephalus sireni</i>	Arkansas	This report
	Illinois	Brooks and Buckner, 1976
	Louisiana	Brooks, 1978
Acanthocephala		
<i>Fessisentis fessus</i>	Illinois	Landewe, 1963; Nickol, 1972; Dunagan and Miller, 1973
	Louisiana	Nickol, 1972; Buckner and Nickol, 1979
<i>Neoechinorhynchus</i> sp.	Illinois	Miller and Dunagan, 1971
Nematoda		
<i>Capillaria</i> sp.	Arkansas	This report
<i>Falcaustra chabaudi</i>	Illinois	Dyer, 1973
	Arkansas	This report
<i>Siren intermedia texana</i>		
Nematoda		
<i>Contracecum</i> sp.	Texas	McAllister and McDaniel, 1992
<i>Siren lacertina</i>		
Trematoda		
<i>Cephalogonimoides sireni</i>	Florida	Premvati, 1969
<i>Diplostomum variabile</i>	*	Harris et al., 1967
<i>Gorgodera minima</i>	Louisiana†	Bennett and Humes, 1938;
	Florida	Brooks and Buckner, 1976
<i>Lechriorchis</i> sp.	Louisiana†	Bennett and Humes, 1938
<i>Progorogodera foliata</i>	Florida	Brooks and Fusco, 1978
<i>Stomatrema guberleti</i>	Florida	Brooks and Buckner, 1976
<i>Telorchis sireni</i>	*	Zeliff, 1937
<i>T. stunkardi</i>	Florida	Brooks and Buckner, 1976
Cestoidea		
<i>Proteocephalus</i> sp.	Louisiana†	Bennett and Humes, 1938;
	Florida	Loftin, 1960
<i>P. aberrans</i>	Florida	Brooks, 1978
Nematoda		
<i>Brevimulticaecum tenuicolle</i>	Louisiana†	Bennett and Humes, 1938
<i>Cosmocercoides dukae</i> or <i>C. variabilis</i>	Florida	Walton, 1938; Baker, 1987
<i>Falcaustra catesbeianae</i>	Florida	Walton, 1938
<i>Siren</i> sp.‡		
Trematoda		
<i>Cephalogonimus amphiumae</i>	Florida	Manter, 1938
<i>Clinostomum marginatum</i>	Florida	Manter, 1938
<i>Diplostomulum</i> sp.	Florida	Manter, 1938

* Data on locality not provided in reference.

† Sirens collected from the campus of Louisiana State University, Baton Rouge, Louisiana, and reported originally to be *S. lacertina*; however, its range does not include Louisiana (Conant and Collins, 1991). In all probability the host is *S. intermedia*.

‡ Species of siren not reported. The ranges of *S. intermedia*, *S. lacertina*, *Pseudobranchius axanthus*, and *P. striatus* include Florida (Conant and Collins, 1991).



Figures 1-4. Larval *Capillaria* sp. infecting the integument of a juvenile *Siren intermedia nettingi* (ASUMZ 16992) from Montgomery County, Arkansas. 1. View of siren showing raised and discolored areas on integument (arrows) infected with larvae. Scale bar = 15 mm. 2. Closer view showing areas of infection on integument. Scale bar = 7.5 mm. 3. Section of skin showing parasite in wall of dermal capillary with severe ectasia of superficial vasculature as well as degeneration and edema in overlying epidermis. Scale bar = 100 μ m. 4. Closer view of parasite in dermal vessel. Note stichostome (arrow). Scale bar = 35 μ m.

1976) Brooks, 1978, were recovered from the duodenum of 9 sirens (173.6 ± 20.7 , 53–252 mm) collected only at the Montgomery County site; mean intensity was 4.3 ± 1.4 (range 1–15) worms. This cestode has been reported previously from *S. i. nettingi* in Illinois and Louisiana (Table 1).

Four kathlaniid nematodes, *Falcaustra chabaudi* Dyer, 1973, were in the rectum of 2 *S. i. nettingi* (235 and 290 mm SVL) collected in February 1991 from Clay County; mean intensity was 2.0 ± 1.0 (range 1–3) worms. This is the first report of *F. chabaudi* from Arkansas. Dyer (1973) described *F. chabaudi* from 2 *S. i. nettingi* in southern Illinois.

A single juvenile *S. i. nettingi* (SVL = 53 mm; ASUMZ 16992) collected in October 1990 from the Montgomery County site was infected with larval *Capillaria* sp. in the integument (Figs. 1, 2). The epidermis of ASUMZ 16992 showed separation of the pigment layer from the underlying basement membrane with moderate hydropic change at the dermal interface. In the dermis there was marked edema with dilation of numerous vascular channels. The larvae adhered to the walls of these dilated vessels (Fig. 3) while provoking no discernable inflammatory response. Dermal connective tissue was unaffected by the presence of this parasite. The stichostome, typical of trichuroid nematodes (Noble et al., 1989), is shown in Figure 4.

A summary of the helminths of *Siren* spp. is presented in Table 1. There appears to be some host specificity among helminths from different siren taxa, particularly among trematodes of lesser and greater sirens. Although *S. intermedia* and *S. lacertina* may be found in sympatry, they appear to be partitioned by habitat differences. The former tends to inhabit more acidic pH waters while the latter is found in aquatic sites with circumneutral pH (P. E. Moler, pers. comm.). This may help explain differences in the trematode faunas of the 2 species as intermediate hosts may also be partitioned in the same manner.

We thank the Arkansas Game and Fish Commission for Scientific Collecting Permits (issued to S.E.T. and B.G.C.), Dr. Steve J. Upton for examining samples for coccidia, and Paul E. Moler for information on sirens.

Literature Cited

- Baker, M. R. 1987. Synopsis of the Nematoda parasitic in amphibians and reptiles. Memorial University of Newfoundland Occasional Papers in Biology 11:1–325.
- Bennett, H. J., and A. G. Humes. 1938. Helminth parasites of *Amphiuma tridactylum* and *Siren lacertina* from Louisiana. Proceedings of the Louisiana Academy of Sciences 4:243–245.
- Brooks, D. R. 1978. Systematic status of protocephalid cestodes from reptiles and amphibians in North America with descriptions of three new species. Proceedings of the Helminthological Society of Washington 45:1–28.
- , and R. L. Buckner. 1976. Some platyhelminth parasites of sirens (Amphibia: Sirenidae) from North America. Journal of Parasitology 62: 906–909.
- , and A. C. Fusco. 1978. Some digenetic trematodes from caudate amphibians in the southeastern United States. Journal of the Mississippi Academy of Sciences 23:95–99.
- Buckner, R. L., and B. B. Nickol. 1979. Geographic and host-related variation among species of *Fessisensis* (Acanthocephala) and confirmation of the *Fessisensis fessus* life cycle. Journal of Parasitology 65:161–166.
- Conant, R., and J. T. Collins. 1991. A Field Guide to Reptiles and Amphibians of Eastern and Central North America, 3rd ed. Houghton Mifflin, Boston. 450 pp.
- Dunagan, T. T., and D. M. Miller. 1973. Some morphological and functional observations on *Fessisensis fessus* Van Cleave (Acanthocephala) from the dwarf salamander, *Siren intermedia* Le Conte. Proceedings of the Helminthological Society of Washington 40:209–216.
- Dyer, W. G. 1973. *Falcaustra chabaudi* sp. n. (Nematoda: Kathlaniidae) from the western lesser siren, *Siren intermedia nettingi* Goin, 1942. Journal of Parasitology 59:994–996.
- Harris, A. H., R. Harkema, and G. C. Miller. 1967. Life history and taxonomy of *Diplostomum variabile* (Chandler, 1932) (Trematoda: Diplostomatidae). Journal of Parasitology 53:577–583.
- Landewe, J. E. 1963. Helminth and arthropod parasites of salamanders from southern Illinois. M.Sc. Thesis, Southern Illinois University, Carbondale, Illinois. 47 pp.
- Loftin, H. 1960. An annotated check-list of the trematodes and cestodes and their vertebrate hosts from northwest Florida. Quarterly Journal of the Florida Academy of Sciences 23:302–314.
- Manter, H. W. 1938. A collection of trematodes from Florida amphibians. Transactions of the American Microscopical Society 57:26–37.
- Martof, B. S. 1973. *Siren intermedia*. Pages 127.1–127.3 in J. D. Anderson, ed. Catalogue of American Amphibians and Reptiles. Society for the Study of Amphibians and Reptiles, American Museum of Natural History, New York.
- McAllister, C. T., and S. J. Upton. 1987. Endoparasites of the smallmouth salamander, *Ambystoma texanum* (Caudata: Ambystomatidae) from Dallas County, Texas. Proceedings of the Helminthological Society of Washington 54:258–261.
- , and V. R. McDaniel. 1992. Occurrence of larval *Contracaecum* sp. (Ascaridida: Anisakidae)

- in Rio Grande lesser sirens, *Siren intermedia texana* (Amphibia: Caudata), from south Texas. Journal of the Helminthological Society of Washington 59:239–240.
- Miller, D. M., and T. T. Dunagan. 1971. Studies on the rostellar hooks of *Macracanthorhynchus hirudinaceus* (Acanthocephala) from swine. Transactions of the American Microscopical Society 90: 329–335.
- Nickol, B. B. 1972. *Fessisentis*, a genus of acanthocephalans parasitic in North American poikilotherms. Journal of Parasitology 58:282–289.
- Noble, E. L., G. A. Noble, G. A. Schad, and A. J. MacInnes. 1989. Parasitology, The Biology of Animal Parasites. Lea & Febiger, Philadelphia. 574 pp.
- Premvati, G. 1969. A new trematode *Cephalogoni-mus sireni* sp. nov. (Digenea: Cephalogoniidae) from Florida mud-eel, *Siren lacertina*. Proceedings of the Helminthological Society of Washington 36:74–76.
- Walton, A. C. 1938. The Nematoda as parasites of Amphibia. IV. Transactions of the American Microscopical Society 57:38–53.
- Zeliff, C. C. 1937. A new species of trematode from the mud-eel (*Siren lacertina*). Proceedings of the United States National Museum 84:223–226.

J. Helminthol. Soc. Wash.
61(2), 1994, pp. 238–240

Research Note

New Host and Distribution Record of *Raillietina* (*Raillietina*) *coreensis* (Cestoda) from *Apodemus argenteus* (Rodentia) in Japan

TAKASHI IWAKI,^{1,3} FRANTISEK TENORA,² AND MASAO KAMIYA¹

¹ Department of Parasitology, Faculty of Veterinary Medicine, Hokkaido University, Sapporo 060, Japan and

² Department of Zoology, University of Agriculture, 613 00 Brno, Czech Republic

³ Present address: Laboratory Animal Research Center, Faculty of Medicine, Tottori University, Yonago 683, Japan

ABSTRACT: *Raillietina* (*Raillietina*) *coreensis* Honda, 1939 is redescribed from the small intestine of *Apodemus argenteus* (Rodentia: Muridae) in Hokkaido, Japan. This report represents a new host and distribution record for *R. coreensis* in the host and in Japan.

KEY WORDS: *Raillietina* (*Raillietina*) *coreensis*, cestode, Davaineidae, *Apodemus argenteus*, rodent, Japan.

Cestodes of the genus *Raillietina* Fuhrmann, 1920 have been reported from various rodents in the tropical and subtropical zones. However, there are few records of *Raillietina* from *Apodemus* Kaup, 1829 (Rodentia). We obtained *Raillietina* (*Raillietina*) *coreensis* Honda, 1939 from a small Japanese field mouse, *Apodemus argenteus* (Temminck) in Hokkaido, Japan, representing a new host and distribution record for the parasite.

Cestodes were collected from the small intestine of an *A. argenteus* (female, 10–18 mo old) captured at Abuta, Hokkaido, Japan in October 1991. Worms were lightly pressed, fixed in 70% ethanol, stained with acetocarmine, dehydrated

in an ethanol series, cleared in xylene, and mounted in MGK® (Matsunami Glass Ind., Ltd., Japan). All measurements in fixed specimens are in micrometers unless otherwise indicated and given as a range with the mean in parentheses.

Raillietina coreensis Honda, 1939 (Figs. 1–3)

Redescription (based on 2 specimens): Total body length 61 and 114 (88) mm, maximum width 2 mm. Scolex 363 and 554 (459) long by 264 and 528 (396) wide. Four suckers oval, 106–123 (117) long by 92–115 (102) wide, with numerous hooks 7–8 long, arranged diagonally with about 7 hooks per row on the inside of the sucker. Rostellum 86–99 (93) wide. Most of rostellar hooks lost during processing. One remaining hook hammer-shaped and 13 long. Proglottids trapezoidal. Mature proglottids 132–238 (180) long by 785–1,465 (1,033) wide. Genital pore unilateral, usually located in anterolateral position in mature proglottids; some situated near middle of margin. Testes 24–29 (27) in number lying on